

Sea surface temperature and heat budget variability in ECCO2


Holger Brix (UCLA)

Dimitris Menemenlis (JPL)

ECCO2 meeting, Boston

23 September 2008

SST anomalies



Mixed layer heat budget and SST

$$\frac{\partial T}{\partial t} = \frac{1}{h} \frac{Q}{\rho C_p} - \frac{1}{h} \frac{\partial h}{\partial t} \Delta T + \text{advection} + \text{diffusion}$$

Project Questions:


- Which processes dominate on which temporal and spatial scales?
- Does spatial averaging matter?
- How big are the associated errors?

Mixed layer heat budget and SST

$$\frac{\partial T}{\partial t} = \frac{1}{h} \frac{Q}{\rho C_p} - \frac{1}{h} \frac{\partial h}{\partial t} \Delta T + \text{advection} + \text{diffusion}$$

Approach:

- Choose three oceanic regions (approx. 5°x5°) in the North Pacific Ocean:
 - the subtropical gyre
 - an upwelling region off the US West Coast
 - a dynamically active area in the Kuroshio region
- Choose different frequency bands:
 - 6 hourly
 - Daily
 - Monthly





SST variability



AVHRR

ECCO2



US West
Coast



North of
Hawaii



East of
Japan





SST anomalies



AVHRR

ECCO2



US West
Coast



North of
Hawaii



East of
Japan



Mixed layer heat budget and SST

$$\frac{\partial T}{\partial t} = \frac{1}{h} \frac{Q}{\rho C_p} - \frac{1}{h} \frac{\partial h}{\partial t} \Delta T + \text{advection} + \text{diffusion}$$

Approach – part 2:

- In a first step calculate $\partial T / \partial t$ and the first two terms on the right hand side – treat advection and diffusion as residual

“Technical” questions:

- Which mixed layer depths?
 - KPPmld vs. KPPhbl
 - Other definitions
- Which ΔT ?